

# Re-assessment of *Arthropleura mammata* from the Moscovian of Northern France: new anatomical information and adaptations to terrestrial environments

Mickaël LHÉRITIER<sup>a,1</sup>, Bruno VALLOIS<sup>b,c</sup>, and Claudie DURAND<sup>b</sup>

The fossil record of *Arthropleura*, distributed in the Nord-Pas-de-Calais coalfield (France) and some Belgian localities, is reassessed and compared to new material discovered in new localities of the Bruay Formation. Based on tergite ornamentation, the Bruay specimens are attributed to *Arthropleura mammata*. Two specimens are preserved with their head, allowing the description of the head anatomy of this species for the first time. We estimated the size of the most complete remains from the newly investigated localities, with a maximum length probably reaching 50 centimetres. The geographical extension of *A. mammata* can be extended to the Nord-Pas-de-Calais and to the Saarland coal basins. Some specimens from the Bruay Formation consist of K-plates, typical ventral plates of arthropleurids displaying well-preserved pierced tubercules. These pierced tubercules are also found in K-plates of *Arthropleura cristata*, found in the Mazon Creek Lagerstätte, and could represent spiracles similar to the ones found in millipedes. This suggests that *Arthropleura* possessed a tracheal system for respiration and, combined with other anatomical factors, would confirm it as a terrestrial arthropod.

Arthropleura | Bruay Formation | Moscovian | Nord-Pas-de-Calais coalfield | K-plates

Received: 03 February 2025; Accepted: 20 May 2025; Published: 01 June 2025

## Introduction

*Arthropleura* Jordan & Von Meyer 1854 has long been an iconic myriapod due to the gigantic size of some of its representatives, that may have reached nearly 3 metres long (Davies et al. 2021), making them the largest land arthropod of all time. *Arthropleura* thrived from the Early Carboniferous (Rößler & Schneider 1997) to the Early Permian (Rößler et al. 2012), with an Euro-American distribution (Hannibal 1997). Their most prominent feature is the division of their tergites between a central syntergite and lateral paratergites. Among members of *Arthropleura*, *A. mammata* Salter 1863 is recognisable by its characteristic tuberculation - a first row of four tubercles under the keel of the paratergite and a second row of smaller tubercles behind the first row. *Arthropleura mammata* was known from the Bashkirian (323.2 Ma – 315.2 Ma) of the United Kingdom, Belgium and France (Salter 1863; Pruvost 1919, 1922, 1930).

In Northern France, knowledge of Arthropleura is largely the result of very limited opportunistic collections, housed in the Natural History Museum of Lille (Hauts-de-France, France). Specimens were found during coal mining at the end of the 19th century and at the beginning of the 20th century (Pruvost 1912, 1919). These fragmentary remains of Arthropleura were examined by Pruvost and already considered as important references for studying the genus Arthropleura. They are deposited under Gosselet collections (acronym MGL, see Appendix 1) and consist of 34 specimens, among which only eight specimens were actually collected during coal exploitation in Nord-Pas-de-Calais localities (Fig. 1). The fossils were determined as Arthropleura mammata (i.e. Pruvost 1912, 1919). Because of a lack of complete specimens, knowledge of this animal has been scarce and characterised by very fragmentary remains examined by Pruvost in 1919. For example none of the examinations include an estimation on the animal size. Based on

previous records, A. mammata has been described as inhabiting the coal marshes of Nord-Pas-de Calais in the locality of Anzin (Pruvost 1912, 1919). This study aims to revisit the biostratigraphy of A. mammata in accordance with plant biozones of Moscovian age, with new material acquired during fieldwork in 1995 to 2012 in the western part of the Nord-Pas-de-Calais coal basin (see Vallois & Nel 2023) (Fig. 1). This material comes from three new localities from the Bruay Formation, forming the upper part of the coal layers. These new occurrences confirm the presence of Arthropleura mammata, already known in the Westphalian B (Duckmantian, late Bashkirian), through Westphalian C (Bolsovian, early to middle Moscovian, see Lucas et al. 2022 for the equivalence between Western Europe and Global stages) and allow a more precise estimation of the size of A. mammata. These new occurrences extend the biostratigraphical span of A. mammata well beyond the Bashkirian age, that would complete a vertical extension of this species (i.e. Fig. 2, red hexagon). A focus is made on samples previously studied by Pruvost (1912, 1919), which correspond to localities in the Valenciennes coal basin of Zeiller 1886 (Anzin Formation). Morphology and ecology of A. mammata is also discussed thanks to the discovery of the first heads observed for this species and ventral plates displaying a possible tracheal system.

Author affiliations: <sup>a</sup>Universite Claude Bernard Lyon 1, CNRS, ENS de Lyon, LGL-TPE UMR 5276, F-69622, Villeurbanne, France; https://orcid.org/0009-0004-0652-2907; <sup>b</sup>Conservation Paléontologie et Paléobotanique, Muséum d'Histoire Naturelle de Grenoble, 1 rue Dolomieu, 38000 Grenoble, France; <sup>c</sup>Levesano (association loi 1901). Association for preservation of historical and cultural patrimonial of Liévin. Maison de la Mémoire, rue du 4 septembre 62800 Liévin, France.

<sup>1</sup> Mickaël LHÉRITIER: micklheritier@hotmail.fr

Managing Editor: Harriet B Drage; Handling Editor: Stephen Pates; Peer reviewers: Anthony Shillito, Carolin Haug, Grzegorz Pacyna; Typesetter:Mickaël Lhéritier





Figure 1. Location of the studied material. A: position of the French and Belgian coalfields. B: position of the Pas-de-Calais basin, where the 'ancient material' (Pruvost, 1919, Anzin Formation, Westphalian B i.e. Baskhirian-Moscovian, green star) and the new specimens from the Bruay Formation (Westphalian C i.e. Moscovian, red hexagon) were found. C: details of the position of the Bruay Formation in the Pas-de-Calais basin. The Bruay specimens come from three different zones exploited by different mining companies: the Bruay-Auchel Group (BAG), the Lens-Liévin Group (LLG) and the Hénin Group (HG). Note that the coalfields are not visible on the map at this scale, being covered by Cretaceous sediments and located in underground mines. An, Anzin; Bb, Borinage basin; Cb, Center basin; Chb, Charleroi basin; FBb, Franco Belgian basin. Scale bars = 10 km (B, C), 500 km (A). Vectorised map in A from © d-maps.com.

# **Geological setting**

The foreland coal basin (covering Northern France and dating from the Moscovian, Fig. 2) is defined by an elongated region of sediment accumulation formed between a compressional orogenic belt and the adjacent craton. The growth of the Variscan mountain range increased sediment supply to the basin due to syn-orogenic erosion, leading to thick sediment accumulations during the Westphalian (Corsin & Corsin 1970). These accumulations are divided into three Formations (locally called Assises): the Vicoigne Formation (*Assise de Vicoigne*), the Anzin Formation (*Assise d'Anzin*) and the Bruay Formation (*Assise de Bruay*). These formations are themselves subdivided into members, following their plant fossil record. Among them, the members of the Bruay Formation are not well differentiated. They are the Six Sillons member, the Ernestine member, the Du-souich member and the Edouard member which are made of a rhythmic succession of parasequences. These sequences include from the base to the top a paleosol, a coal seam, shales rich in plant fossils and conglomerates and sandstones both with a decreasing sorting (Bouroz 1969; Dollé 1970; Le Gall 1994).

The Bruay Formation, where *Arthropleura* was most abundant, is separated from the preceding Formation (Anzin) by the Rimbert marine band (Fig. 2) indicating the last marine transgression. The Bruay Formation includes a repetition of parasequences, which enabled the formation of 37 coal seams with a thickness varying from 770 to 1490 metres (Corsin & Corsin 1970). These parasequences are an indicator of a subsidence directed by climatic, eustatic and tectonic conditions. Within each sequence the organic deposit responsible for the coal seams accumulated in a calm, shallow and anoxic environment similar to modern bogs or mangroves. These seams were covered by sedimentary material with decreasing sorting. This sedimentary material is composed from the bottom





Figure 2. Biostratigraphical ranges of key Arthropleura species from northern France basins (including Belgian occurrences). Correlation with plant biozone markers from Zieller (1894) and faunistic biozone markers from Hennion et al. (2015) and Schneider et al. (2021). Each sequence of formation, successively Vicoigne, Anzin, Bruay, refers to eponymous localities and is adapted from the stratigraphic column figured in Corsin & Corsin (1970). Green star: eight old historical references (Pruvost, 1919, Anzin Formation). Red hexagon: Bruay Formation.

to the top of shales deposited during the transgressive context of a deltaic plain, then flooding plain sandstones deposited in a more regressive context.

Zeiller (1894) was among the first to use fossil plants for stratigraphy in these basins. His monography on the Carboniferous of the Valenciennes basin (Zeiller 1886) introduced a new macrofloral approach, that successfully correlated formations based on the plant fossils abundance and similarity of plant sequences, see in Figure 2 (B3, C1, C2 subzones). The productive coal seams in the subzones B1 and B3, exploited in the Anzin area, led to the designation of Anzin as the type locality of the Anzin Formation. This formation is considered a typical series of Westphalian B (Bertrand 1914), equivalent to the late Bashkirian in international time scale. These coal seams are located in underground mines and not in outcrops.

Although the Bruay-les-Mines locality and adjacent areas (e.g., Lens-Liévin and Hénin-Liétard coal districts, Pas-de-Calais, France) have been exploited since the 19th century, the Bruay Formation was defined more recently (Corsin & Corsin 1970; Vallois & Lebrun 2010). This formation includes the C1 and C2 subzones of Zeiller (Fig. 2). These plant sequences were later identified as belonging to the Westphalian C or the equivalent Early Moscovian. In the Nord-Pas-de-Calais coal basin, correlations between different formations and members are based on plant remains and tonsteins (Bouroz et al. 1963; Bouroz 1967), which are fine deposits, often of volcanic origin, which could cover large areas (grey vertical boundaries in Figure 2).

# Materials and methods

The studied material comes from the Anzin and Bruay Formations. Specimens from the Anzin Formation mainly come from outcrops in the vicinity of the Valenciennes area. This material was described in previous literature (Pruvost 1912, 1914, 1919, 1930), that we refer to as "ancient material" (green star in Figs 1 and 2) and is housed in Lille Museum (Gosselet Collections, acronym = MGL). Reassessment of Lille specimens made by previous authors (Pruvost 1912, 1919) and by the curator of Gosselet collections (Oudoire, pers. comms.) gave a clear view on the disparity of *Arthropleura* (see Appendix 1 for the re-assessment of this material and Appendix 2 for additional data on Lille collections). Lille specimens consist



of leg fragments (MGL.1874 and MGL.5037-3, Pruvost 1912), ventral plates (MGL.1877, Pruvost 1919), a syntergite fragment (MGL.1878a-b, Pruvost 1919), another syntergite fragment associated with finely wrinkled soft parts (MGL.1881a-b, Pruvost 1919) and a paratergite fragment (MGL.5057-1). Specimens from the Bruay Formation were collected from three localities between 1995 and 2019:

- Slag heap of pit N° 5 of Bruay-Auchel group (acronym: BAG -50° 28' 49" N, 2°31' 10" E )
- Slag heap at pit N° 4 East (50° 24' 48" N, 2° 59' 48" E) and N° 3 East of Henin Group (acronym: HG - 50° 24' 46" N, 2° 56' 22" E)
- Slag heap at pit N°7 of Lens-Liévin Group (acronym LL 50° 24' 09.98" N, 2°48' 50.61" E)

This newly investigated material (red hexagon in Figs 1 and 2) is housed at the Museum of Natural History of Grenoble (Isère, France), under label MHNGr.PA.

# Systematic palaeontology

Subphylum **Myriapoda** Latreille, 1802 Total-group **Diplopoda** de Blainville in Gervais, 1844 or total-group **Pectinopoda** Benavides, Edgecombe and Giribet, 2023

**Remarks**. This systematic position is based on the position of *Arthropleura* obtained in the latest total-evidence analyses from Lhéritier et al. (2024).

#### Subclass Arthropleuridea Waterlot, 1934

**Diagnosis** (emended from Edgecombe in Minelli, 2015). Myriapoda with diplosegmentation and paired anal valves. Paratergal lobes separated from the syntergal axis by sutures. Sclerotized plates (K, B and rosette plates) buttressing the leg.

#### Order Arthropleurida Waterlot, 1934

**Diagnosis** (modified from Wilson, 1999 and Edgecombe in Minelli, 2015 and adapted from the observations in Lhéritier et al. 2024). Arthropleuridea with ventral sclerites at the front of the head. Eyes stalked. Antennae with eight articles. Mandible with gnathal lobe separated from the distal part of the mandible base. First maxillae reduced with short palp. Second maxillae present. At least 23 trilobed dorsal plates bearing irregular large and small tubercles. Two sternites per dorsal tergite. Sternites with a rounded posterior margin and a more pointed anterior margin. Groove crossing the sternites midline.

**Remarks**. Contrary to what was said in Lhéritier et al. 2024, the antennae of *Arthropleura* (notably those observed in Montceau-les-Mines juveniles) bear eight articles (not seven).

#### Family Arthropleuridae Von Zittel, 1885

**Diagnosis**. Same as order. **Remarks**. Monogeneric family.

Genus Arthropleura Jordan in Jordan and Von Meyer, 1854

**Type species.** Arthropleura armata Jordan 1854. Holotype  $K_{13-14}$  deposited in the Museum für Naturkunde, Berlin. **Diagnosis.** Same as family.

#### Arthropleura mammata Salter 1863

1912 Arthropleura mammata Pruvost: 57-64, pl.II figs 1-3, 5 1919 Arthropleura mammata Pruvost: 81, pl. XXV figs 7-8, 10, 20

**Type material**. Lectotype UKBGS.GSM 25381 deposited in the British Geological Survey.

**Type stratum and locality**. Upper Carboniferous, Westphalian B, top of the *Modiolaris* Zone in the Ferny Metal, beneath the Seven Foot Coal of Rams Mine; Pendleton Colliery, near Manchester, Lancashire, UK.

Referred Material. MHNGr.PA.39321 (Figs 3A, B and 4). MHNGr.PA.39322 (Fig. 5A) MHNGr.PA.39324 (Fig. 5C) MHNGr.PA.39327 (Fig. 6F) MHNGr.PA.39328-1 (Fig. 5B) MHNGr.PA.39347 (Fig. 6E) MHNGr.PA.39551-1 (Fig. 7A, C) MHNGr.PA.39551-11 (Fig. 7B, D) MGL.1878 (Fig. 6A, B) MGL.5057-1 (Fig. 6C) MGL.5057-3 (Fig. 6D)

**Origin of the material**. Slag heap N°1A, pit N° 5 of Bruay at Bruay-Auchel group (50° 28' 49" N, 2° 31' 10" E ), near Divion, Pas-de-Calais- France. See the Material and methods section for the other pits.

**Age and stratigraphy**. Early Moscovian, Westphalian C or Bolsovian (Substages Western Europa) – Bruay Formation.

**Diagnosis** (modified after Salter 1863). *Arthropleura* with round and heavily ornamented head. Collum without paratergites. Trunk paratergites bearing very granulated furrow and no tubercle behind the keel. Posterior part of paratergites with one row of large and one row of smaller tubercles. Middle to front part of the syntergite granulated.

**Description**. Head rounded anteriorly, ranging in length from 23.2-33.8 mm and in width from 41.7-50 mm, and bearing 50 (MHNGr.PA.39321, Fig. 3AB) to 61 (MHNGr.PA.39322, Fig. 5A) tubercles (1.1-6.4 mm in width). Tubercles round with a diameter of 1.4 to 5.4 mm. Number of tubercles decreasing from the anterior to the posterior part. Ventral sclerites (maximum width of 4.9 mm in MHNGr.PA.39322) at the front of the head (Fig. 5A) in the shape of flattened plates beginning on the sides of the head and rejoining at the centre. Ventral sclerites wider laterally and decreasing in width at the centre.

Specimen MHNGr.PA.39321, the most complete, comprises nine body segments (Fig. 3). First segment modified into a collum (13.1 mm long and 41.7 mm wide) and different from the following segments by being composed of only a syntergite (Figs 3 and 4). Segments two to nine divided into a central syntergite and lateral paratergites (Fig. 3). Average syntergite length and width: 14.5 and 39.4 mm, respectively. Middle to front part of the syntergite granulated (Figs 3, 4, 5). Only left paratergite (37.2 mm wide and 28.6 mm long) of segment five observed (Fig. 3). Paratergite with a row of two visible tubercles (diameter 5 to 7 mm) (Figs 3, 4, 6E) followed by a second row of smaller tubercles (diameter 4,4 to 3 mm) (Figs 3, 4, 6E).

MHNGr.PA.39551-5 and MHNGr.PA.39551-11 are K-plates. These plates are 10 mm long and 5 mm (MHNGr.PA.39551-5) (Fig. 7A, C) to 6 mm (MHNGr.PA.39551-11) (Fig. 7B, D) wide. They are divided into five parts (Fig. 7). The first one (more proximal part to the leg) is triangular and the largest one, with a maximum length and width of 6 mm and 5 mm respectively. It is elongated toward the





Figure 3. Arthropleura mammata, specimen MHNGr.PA.39321. A: whole specimen. B: detail of the head. C = collum; H = head; Pt = paratergite; St = syntergite; T = tubercule; T# = tergite number #. White rectangle on A corresponds to the position of B. Scale bars = 10 mm (B), 20 mm (A). Photos by C. Durand, Muséum d'Histoire Naturelle de Grenoble.





Figure 4. Reconstruction of Arthropleura mammata. Based on MHNGr.PA.39321. C, collum; H, head; Pt, paratergite; S, syntergite. Numbers indicate the segments observable on the specimen. Scale bar = 120 mm.



leg (Fig. 7). The second part (middle of the plate, 3 mm in length and 5 mm in width), is also triangular pointing toward the sternite (Fig. 7). The third part (most distal part to the leg and closer to the sternite), is sub-triangular (2 mm long and 5 mm wide; Fig. 7). The fourth part (behind the third part) is sub-triangular (2 mm long and 1 mm wide), pointing toward part three (Fig. 7B, D), and seems to be located at the boundary between the K-plate and the sternite. The fifth part (only observed in MHNGr.PA.39551-11), is anterior to the first and second parts (Fig. 7). It is trapezoidal (length of 5 mm and a width of 1 mm) and seems to be located at the boundary between the K-plate and the coxa (B-plate) of the next leg in front. Multiple pierced tubercles are present on all parts of the plates (from 91 to 103 on the first part, from 32 to 38 on the second part, 21 on the third part, eight on the fourth part and 12 on the fifth part) (Fig. 7). Tubercle roughly round with a diameter of 0.2 to 0.63 mm. Opening of the tubercle also round with a diameter of 0.07 mm.

# **Discussion**

### **Extent of Ancient specimens**

Several fragments of Arthropleura mammata have been discovered in the past, the stratigraphical extent of these ancient specimens is represented by a green star in Figure 2. Most were found near the eponymous locality of the Anzin Formation, Anzin. The discovery of these specimens seems to be linked to the historical context of coal mining in this region during the 19th and 20th century. Arthropleura mammata was considered as living in a spatially restricted zone with occurrence in the fossil record also restricted stratigraphically. These coal exploitations seemed to be limited geographically near Anzin and Valenciennes localities (Nord Department, Hauts-de-France region) and in excavation dumps, comprising material from the type locality of the Anzin Formation. The coal exploitations are mainly located in the eastern part of the Nord-Pas-de-Calais coal basin. Dated from the late Bashkirian (Westphalian A and B), these deposits are located between "La Passée de Laure marine band" and the "Rimbert marine band", corresponding to a time interval in which a rich flora arose (Molina-Solís et al. 2024), qualified by Zeiller as the "Valenciennes basin" flora (Zeiller 1886) (Fig. 2). However, It should be noted that A. mammata discoveries remained rare even during coal mining times, during which the majority of A. mammata specimens near the locality of Anzin were found (Pruvost 1912, 1919).

#### New extent in Bruay Formation

The new occurrences from Bruay are younger than previous occurrences of *A. mammata* found in Northern France (Anzin locality), extending the range of this species in the Nord-pas-de-Calais coal basin from the Bashkirian to the early Moscovian (Westphalian C). Its stratigraphy could possibly extend also into the late Moscovian (Westphalian D), based on the macroflora assemblage observed in the excavation dump n°76 from Liévin (Fig. 6C). Among the Bruay occurrences, specimen MNHGr.PA.39321 (Fig. 3) is among the most complete specimens of *Arthropleura mammata* found yet and can be compared to specimen MTB 1220 of *A. armata* found in equivalent chronostratigraphic ranges in the Saar basin (Brauckmann et al. 1997; Wilson 1999).

#### **Congruent extents in European localities**

A. mammata, like other arthropleurid species, is mostly known in continental paleo-equatorial environments (Schneider et al.

2010). The stratigraphical extension of A. mammata goes from the Bashkirian (Pruvost 1912, 1919, 1930) to the early Moscovian (new Bruay occurrences). We propose that its extension was uninterrupted from the Bashkirian to the Moscovian even through the Carboniferous rainforest collapse, recorded by four major marine transgressions (Fig. 2, Corsin & Corsin 1970; Molina-Solís et al. 2024). The oldest occurrences of the genus Arthropleura are known from the Visean in the Hainichen basin in Germany (Rößler & Schneider 1997). No Arthropleura occurrence from this age is found in the Nord-Pas-de-Calais basin (acronym HBNPC), which during this stage was a marine environment with a scarce Namurian continental flora assemblage. In the lower part of the HBNPC, the marine nature of the deposits is confirmed by the presence of Goniatites (Chalard 1960). Some rare continental outcrops yield remains of *Pecopteris aspera* Brongniart 1828, indicating the existence of some islands bearing flora at this time.

Among older records, occurrences are found during the Serpukhovian in the Lower and Upper Silesian basins in Poland (Pacyna et al. 2018) and a large specimen of Arthropleura was described in the Stainmore Formation from the Serpukhovian of England (Davies et al. 2021). Disparate elements of Arthropleura have been described in the Bashkirian in the Ostrava-Karvina basin in Czechia (Řehoř & Řehořova 1972), the Charleroi Formation in Belgium (Pruvost 1930), the Donets Basin in Ukraine (Dernov 2019), the Pennine Lower and Middle Coal Measures Formation in the United Kingdom (Vernon 1912; Andrée 1913; Pruvost 1930; Anderson et al. 1997), the Lower Silesian Basin (Zacler Formation) and the Upper Silesian Basin (Ruda Beds) in Poland (Pacyna et al. 2012; Pavela 2018) and the Emma Mine in the Netherlands (Pruvost 1930). Localities increase in abundance in the upper part of the HBNPC with a maximum of recorded occurrences in the Moscovian with specimens in Germany (Jordan & Meyer 1854; Andrée 1913; Guthörl 1934, 1935; Hahn et al. 1986; Kraus & Brauckmann 2003; Kraus 2005), the United Kingdom (Salter 1863; Woodward 1907; Andrée 1913; Pruvost 1930; Hahn et al. 1986; Proctor 1998), France (Waterlot 1934), the Netherlands (Pruvost 1930), Sardinia (Pillola & Zoboli 2021) and in Czech Republic (Pavela 2018). This rise in recording is not simply correlated with the generalised coal mining in localities from this age in Europe and North America and the associated sampling bias. This peak of Arthropleura occurrences is not in contradiction with the new described material. 105 fragments of Arthropleura from the early Moscovian were already described in the Saar basin (Waterlot 1934; Guthörl 1936). The extension of A. mammata seems to be in congruence with a peak of macrofloral diversity that characterises the wet environments of Variscan foreland basins (Molina-Solís et al. 2024). Recently, Pillola & Zoboli (2021) described a paratergite of Arthropleura armata in Sardinia (Italy). This occurrence is the most meridional one found so far, and confirms a late Moscovian extension for the genus.

#### **Chronostratigraphic implications**

Until now, most *Arthropleura* specimens found in the HBNPC were grouped under the species *Arthropleura mammata*, based on eight isolated elements (MGL collection), found solely in the Anzin Formation (Pruvost 1912, 1914, 1919, 1930). The stratigraphical extent of *A. mammata* is thoroughly described in multiple monographs and theses (Brongniart 1828; Boulay 1876, 1879; Zeiller 1886; Carpentier 1913; Danzé-Corsin 1953; Danzé 1956; Buisine 1961; Crookal 1964, 1969; Laveine 1967; Laveine et al. 1978, 1987, 1992; Laveine & Goubet 1995; Izart et al. 1998; Laveine & Belhis 2007).





Figure 5. Arthropleura mammata, other specimens from La Bruay Formation. A: head, specimen MHNGr.PA.39322. B: syntergite fragment, specimen MHNGr.PA.39328-1. C: paratergite fragment, specimen MHNGr.PA.39324. C = collum; T = tubercule; Vs = ventral sclerites. White arrow in C indicates an *Alethopteris serlii* pinnule. Scale bars = 10 mm. Photos from C. Durand, Museum d'Histoire Naturelle de Grenoble.





Figure 6. Arthropleura mammata, specimens from the Anzin and Bruay formations. A-D: Anzin Formation. A-B: part and counterpart of a syntergite, specimen MGL.1878 (same as in Pruvost, 1919, pl. XXV, fig. 8). C: paratergite, specimen MGL.5057-1, Vicq locality, Pit Cuvinot, Westphalian B, Meunière member, seam Boulangère. D: fragment of leg segment, specimen MGL.5057-3, same locality as MGL.5057-1. E-F: Bruay Formation, Lens-Lievin Group. E: paratergite, specimen MHNGr.PA.39327. F: fragments of paratergite and syntergites, specimen MHNGr.PA.39347. White arrows on C and F indicate a *Linopteris subbrongniartii* pinnule. Scale bars = 10 mm. E and F made by C. Durand, Muséum d'Histoire Grenoble.





Figure 7. Arthropleura mammata, K-plates. A: right K-plate, specimen MHNGr.PA.39551-5. B: left K-plate, specimen MHNGr.PA.39551-11. C: details of the central part of MHNGr.PA.39551-5. D: details of the central part of MHNGr.PA.39551-11. 1 = part proximal to the leg; 2 = middle part; 3 = part most distal to the leg; 4 = connexion between K-plate and B-plate. Small white arrows indicate the orientation of the plate: I = leg; s = sternite. White arrows on C and D indicate pierced tubercles. Scale bars = 2 mm (C, D); 5 mm (A, B).

Plant remains associated with *Arthropleura* specimens known in the Anzin Formation are also present in the Bruay Formation (Vallois & Lebrun 2010; Vallois & Nel 2023; Molina-Solís et al. 2024). Some seed fern specimens can be identified (Bertrand 1907), indicating in the Bruay Formation a *Paripteris linguaefolia* (Bertrand 1930) biozone (Fig. 2). It seems that the increase of macroflora diversity is synchronous with the increase in *Arthropleura* occurrences from the Bashkirian to the late Moscovian.

The successive Westphalian series characterising the HBNPC in the Vicoigne, Anzin and Bruay Formations are not very differentiated. They are formed of a rhythmic succession of parasequences, consisting from the base to the top of a paleosol, a coal seam, shales rich in plant fossils and conglomerates and sandstones with a decreasing sorting (Bouroz 1969; Dollé 1970). The repetition of these sedimentary sequences highlights rapid changes of the hydrodynamic conditions, which have been controlled by variations of climatic, eustatic (Fig. 2, red lines indicating marine transgressions), and tectonic conditions. The climax of these variations was reached during the Asturian (Westphalian D), identified in the HBNPC by the Du-souich and Edouard members (Fig. 2). With each cycle, the organic matter which produced the coal seams accumulated below stagnant water or a shallow bog or mangrove, forming an anoxic environment. This organic matter, including possible *Arthropleura*  exuviae was covered by transgressive clay (becoming shales with diagenesis) from deltaic plains, then by fluviatile sandstones from flood plains, which have been deposited in a regressive context.

As seen in Molina-Solís et al. 2024, the specific richness of plants increases exponentially and corroborates with A. mammata occurrences in the Bruay Formation. We disagree with the idea that an Arthropleura association was strictly limited to the Valenciennes basin in the Anzin-Valenciennes basin during the Bashkirian like said in ancient studies (Pruvost 1912, 1919, 1930). The occurrences in the Bruay-Auchel mining group extend the stratigraphy of Arthropleura. This extension affirms that Arthropleura was present along the whole stratigraphical sequence of the basin from the Bashkirian to the Moscovian nearly without interruption. The remains of Arthropleura have been found starting from the Olympe member (Fig. 2), where plant richness reaches around 60 species at the base of the Vicoigne Formation (marked by Lyginopteris hoeninghausii Potonié 1897, Fig. 2). The swampy environment of this time is characterised by an increase of pteridosperms and non-marattialean ferns, observed with occurrences on specimens MGL.1878 and MGL.1881 (Figs 2 and 6A-D). The diversity of plants still increased progressively in the Vicoigne and Anzin Formations (Langsettian-Duckmantian i.e. Westphalian A-B or late Bashkirian), reaching around a hundred species in the late Westphalian B, as





Figure 8. Arthropleura cristata, Mazon Creek, Moscovian-Kasimovian and reconstitution of the tracheal system. A-C: specimen FMNH PE9304. A: part. B: counterpart. C: details on the pits. D: unnumbered specimen (named specimen FMNH 1 in Wilson 1999). E-F: reconstitution of the tracheal system. E: position of the supposed tracheal system (blue) on the K-plate. F: Details on the K-plate, highlighting the pierced tubercules as possible tracheal openings. Square in A indicates the position of C. 1 = part most proximal to the leg; 2 = middle part; 3 = part most distal to the leg; 4 = connexion between K-plate and sternite; 5 = connexion between K-plate and B-plate. B-P = B-plate; K-P = K-plate; St = sternite; T = pierced tubercule. White arrow on C indicates pierced tubercules. Scale bars = 1 cm (C); 2 cm (A, B). A, B and C taken by Mane Pritza and provided by Paul Meyer. D from Wilson 1999.

seen in the Pouilleuse member (Fig. 2), ending with the marine transgression of the "Rimbert Marine band".

The floral diversity seems in accordance with *Arthropleura* ecology, as a detritivorous and/or possibly herbivorous (Lhéritier et al. 2024) myriapod feeding on abundant plant biomass available in swamp areas in Northern France (Rolfe 1985) (see Appendix 2 for additional knowledge of presumed *Arthropleura* diet). A new increase of plant diversity is recorded in the lower part of the Bruay Formation (Bolsovian i.e. Westphalian C or early Moscovian). This occurs starting from the floristic zone B3 *sensu* Zeiller 1896 (Molina-Solís et al. 2024, Fig. 2). A diversity peak is observed in the upper part of the Six-Sillons member (> 110 plant species), including a high number of non-marattialean ferns and medullosalean pteridosperms. This interval is correlated with the presence of *A. armata* (specimen MGL.1877), considered here as

a preliminary occurrence from the upper part of the member but also from the same age as *A. mammata* specimens from the Bruay Formation.

#### Size and ontogeny

The animal is reconstructed in Figure 4, on the basis of *Arthropleura armata* from Sulzbach (Saarland, Germany; Moscovian; Davies et al. 2021), it is estimated to be 50 centimetres long. The old specimens from the Anzin Formation in the Gosselet collection of the Lille Museum seem to indicate that there were already large specimens (probably adult fragments) since the Bashkirian. Comparable large *Arthropleura* have been assigned to *A. mailleuxi* Pruvost 1930, but the lack of preserved appendages, head, or detailed ornamentation in the *A. mailleuxi* specimens precludes confident species-level



comparisons. Specimens MHNGr.PA.39321 and MHNGr.PA.39322 document the first complete head cuticle of adult Arthropleura mammata. Complete heads are only known in Montceau juvenile specimens (Lhéritier et al. 2024), only fragments are preserved in other localities (Kraus & Brauckmann 2003). The head of Arthropleura seems to keep the same shape during ontogeny, as it is the case for extant centipedes and millipedes that are its closest relatives (Lhéritier et al. 2024). In these two groups, there is no drastic morphological change of the head during ontogeny, the major change being an increase in size (Minelli 2011, 2015). The presence of ventral sclerites at the front of specimen MHNGr.PA.39322 (Fig. 5A), also observed in Montceau juveniles (Lhéritier et al. 2024), confirms that these structures are persistent throughout ontogeny. No other appendages (antennae, feeding apparatus or eyes) were observed in the Bruay specimens. A. mammata head morphology is very close to that of Montceau specimens and may have displayed comparable head appendages (i.e., stalked eyes, antennae, mandibles and maxillae; Lhéritier et al. 2024). A. mammata could have followed an hemianamorphic development (i.e., the addition of new segments at each mould until the sexual maturity followed by multiple moulds with an increase in size; Minelli 2015) like seen in the Montceau juveniles. Ontogenetic development is quite stable among myriapod clades (e.g., all millipedes follow an anamorphic ontogeny), hence it is plausible that all arthropleurids followed the same developmental pattern. As the posterior part of MHNGr.PA.39321 and MHNGr.PA.39322 are not preserved, we cannot see if the posterior segment and legs were not fully developed as in juveniles (Lhéritier et al. 2024) and assess if the specimens were fully matured or juveniles.

#### Habitat

The association of Arthropleura species with plant remains of the Lonchopteris rugosa Brongniart 1822 zone (e.g., Alethopteris davreuxii (Brongniart 1836), Mariopteris muricata (Brongniart), Sphenophyllum miriophyllum (Crepin)) (Bouroz 1940 and Corsin & Corsin 1970) suggest that these Arthropleura inhabited wetlands during the Bashkirian. The presence of Paripteris pseudogigantea (Potonié) Gothan 1953, P. linguaefolia and Laveineopteris rarinervis (Bunbury) Cleal et al., 1990, typically found in early Moscovian (Bolsovian) deposits, may indicate environmental changes in Arthropleura's habitat. In the western part of the HBNPC, no Arthropleura specimens are known prior to the Moscovian surely because it was a marine environment as indicated by the Rimbert marine band (Fig. 2). It is only after this marine band that we observe Arthropleura along with the increase of neuropterid plants occurrences (Corsin & Corsin 1970; Vallois & Nel 2023). Arthropleura was also recovered in Belgium, in coal series that developed between the Quaregon Marine Band and the Petit-Buisson Marine Band (Westphalian A and B; Pruvost 1930; Paproth et al. 1983). The occurrences of Arthropleura mammata in the two coalfield series, recorded in the French and Belgian sites, are dated of the same time interval. Due to the occurrence of Arthropleura in both series before and after the submergence by the sea, we consider that the geographical extension of Arthropleura may have been reduced due to environmental changes, limiting its presence to the coalfields. This new discovery of *A. mammata* in the Bruay Formation helps to build a better understanding of the Carboniferous terrestrial fauna of the HBNPC. The scarcity of animal remains in the Carboniferous of Northern France could be linked to these dynamic changes of their habitats (Molina-Solís et al. 2024). Taphonomic parameters could also explain this stratigraphic pattern, as coalfields are the most suitable Carboniferous environments for preserving *Arthropleura* and other fauna remains, compared to other drier environments.

#### Respiration

All extant myriapods are terrestrial and breathe through a tracheal system or cuticle diffusion (Minelli 2011, 2015). The respiratory system of Arthropleura is still debated as there is still no fossil evidence of spiracles (tracheal openings). The K-plates, observed here in MHNGr.PA.39551-5 and MHNGr.PA.39551-11, are usually interpreted as being associated with possible tracheal pouches based on the fact that the so-called "K-plates" are in fact not plates but bag-like structures (Wilson 1999; Kraus & Brauckmann 2003). Multiple specimens of Arthropleura cristata Richardson 1959 from the Kasimovian Mazon Creek Lagerstätte show the same pierced tubercles on the K-plates (Fig. 8A-C) and one shows internal details of the K-plates (Wilson 1999), especially a network of reticulated structures composed of small interconnected chambers and circular cavities (Wilson 1999) (Fig. 8D). In this context, the tubercles with openings observed in our specimens may represent the outer surface of such cavities (Fig. 8E, F). One major counter-argument to the presence of a tracheal system in Arthropleura is the lack of spiracles observed on the sternites (where they are located in extant millipedes, Minelli 2015). However, Wesener et al. 2014 discovered that the spiracles of glomeridesmids are not present on the sternites but on another plate they called the "stigmatic plate" which is connected to the coxa. As Arthropleura is a stem-relative of Diplopoda (i.e. millipedes; Lhéritier et al. 2024), the spiracles, if present in Arthropleura may not be located on the sternites but on another plate equivalent to the stigmatic plate (in this case the K-plate). Even if a tracheal system can also be found in aquatic arthropods like aquatic insects, the general anatomy of Arthropleura is really similar to extant myriapods (that are all terrestrial) and with no clear adaptations to life in water (modified tracheae, swimming setae on legs). This would imply that Arthropleura was not an aquatic but a truly terrestrial arthropod.

## Conclusions

Arthropleura mammata from the Bruay Formation (early Moscovian) reveals important information of the head and ventral K-plates of this species' adult specimens. This new material extends the stratigraphic range of A. mammata to the Moscovian. The presence of the same group of plants associated with A. mammata throughout its stratigraphic range and its disappearance in the fossil record in intervals with sea submergence could suggest that A. mammata preferentially lived in low altitude wetlands. The heads of the Bruay specimens are also among the most complete adult heads of the genus, and can be compared to the better-preserved ones of Montceau-les-Mines juvenile specimens. The presence of ventral head sclerites (previously only seen in Montceau specimens) in adult representatives confirm their persistence throughout their development. The presence of multiple pierced tubercles on Kplates suggests that Arthropleura had a tracheal system, with the holes being its openings towards the exterior surroundings. This interpretation is supported by comparable features found in Mazon Creek Arthropleura cristata specimens displaying the same pierced tubercles on their K-plates. If the respiratory function of these structures is confirmed, then Arthropleura is likely to have been an air breathing terrestrial arthropod, as all extant myriapods. The Mazon Creek Lagerstätte bears interesting material for the detailed



study of the ventral plates and the possible 3D reconstruction of the respiratory system of *Arthropleura*.

## **Data availability**

Specimens of *Arthropleura mammata* are deposited in the Lille Museum and the Museum of Natural History of Grenoble. Specimens from the Bruay Formation can be found in an online database (https://paleo-collections.museum-grenoble.fr/fr/collections/arthropleura-collection-museum-

grenoble). Specimens of *Arthropleura cristata* figured in Figure 8A-C are deposited in the Field Natural History Museum of Chicago. All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary data.

# Acknowledgements

We would like to thank particularly Hervé Duquesne, generous donator of studied material, and deceased Louis Marie Legrand, our tireless colleagues in the field excursion on slag heaps of Northern France. Missed Emeritus Professor Jean-Pierre Laveine, deceased, for the kind authorization he obtained at that time from the operator Dufour Society for access to the different slag heaps examined. We also thank Thierry Oudoire, previous curator in charge of the palaeontological collection, for access to the historical collection of the Museum of Natural History of Lille, Jessie Cuvelier in charge of the collections at the university of Villeneuve d'Ascq, Rebecca Bilon, director of the Grenoble Natural History Museum and Daniel Verhelst for its Arthropleura reconstruction at the museum. We thank Mane Pritza and Paul Mayer (Field Natural History Museum of Chicago) for providing pictures of Arthropleura cristata. We thank the editor Stephen Pates and our three reviewers Anthony Shillito, Carolin Haug and Grzegorz Pacyna for their constructive advice and editorial assistance that helped improving the quality of our manuscript.

## **Author contributions**

M.L. and B.V. contributed to the design of this study. B.V. and C.D. provided the specimens for this study. B.V. and C.D took photos of the specimens. M.L and B.V. conducted the preparation and improvement of data visualisation. All authors prepared the original draft.

## **Competing interests**

The authors declare no competing interests.

#### Ethics and Al use

No AI was used for this study.

# References

ANDERSON, L. I., DUNLOP, J. A., HORROCKS, C. A., WINKELMANN, H. M. and EAGAR, R. M. C. 1997. Exceptionally preserved fossils from Bickershaw, Lancashire UK (Upper Carboniferous, Westphalian A (Langsettian)). *Geological Journal*, 32, 197–210.

ANDRÉE, K. 1913. Weiteres über das carbonische Arthrostraken-Genus Arthropleura Jordan. *Palaeontographica (1846-1933)*, 295–310.

BERTRAND, P. 1907. Végétaux houillers de Bruay. Ann. Soc. Geol. Nord-FR, 36, 141–145. ———. 1914. Les zones végétales du terrain houiller du Nord de la France. Leur extension verticale par rapport aux horizons marins. *Annales de la Société Géologique du Nord*, 43, 208–254.

BOULAY, N. 1876. Le terrain houiller du Nord de la France et ses végétaux fossiles. Lefebvre-Ducrocq.

——. 1879. Recherches de paléontologie végétale dans le terrain houiller du nord de la France (concession de Bully-Grenay). F. Hayez.

BOUROZ, A. 1940. 1re thèse: Facies et massifs de végétation dans la formation Houillère du Nord de la France: application aux travaux de recherche miniers.Université de Lille, 240pp.

——. 1967. Correlations des tonsteins d'origine volcanique entre les bassins houillers de Sarre-Lorraine et du Nord-Pas-de-Calais. *Comptes Rendus de l'Académie des Sciences de Paris.* 

——. 1969. Le Carbonifère du Nord de la France. *Soc. Geol. Nord, Ann*, 89, 47–65.

——, STIEVENARD, H., BUISINE, A., DALINVAL, A., DOLLE, P., PINNEL, G. and PUIRABORD, G. 1963. Carte des zones stratigraphiques à la côte stratigraphique—300 m du bassin houiller du Nord et du Pas-de-Calais. Service de Ressources et Etudes Géologiques des HBNPC, Institut Géographique National, Paris.

BRAUCKMANN, C., GRÖNING, E. and THIELE-BOURCIER, M. 1997. Kopfund Schwanz-Region von *Arthropleura armata* Jordan, 1854 (Arthropoda; Ober-Karbon). *Geologica et Palaeontologica*, 31, 79–92.

BRONGNIART, A. 1828. *Prodrome d'une histoire des végétaux fossiles*. Levrault.

BUISINE, A. 1961. Contribution à l'étude de la flore du terrain houiller. Les alethopteridées du Nord de la France. Études géologiques pour l'Atlas de topographie souterraine. I. Flore fossile.

CARPENTIER, A. 1913. Contribution à l'étude du Carbonifère du Nord de la France.Université de Lille.

CHALARD, J. 1960. Les horizons à goniatites du Namurien du Nord de la France. *CR IVe Congrès pour l'avancement des Etudes de Stratigraphie et de Géologie du Carbonifère, Heerlen*, 1, 87–92.

CORSIN, P. and CORSIN, P. 1970. Un siècle et demi de Paléobotanique dans le Nord de la France. *Annales de la Société Géologique du Nord*, 90, 223–252.

CROOKAL, R. 1964. Fossil Plants of the Carboniferous rocks of Great Britain. [Second section]. *Mem. Geol. Surv. Great Britain Palaeont.*, 4, 217–354.

——. 1969. Fossil Plants of the Carboniferous rocks of Great Britain. [Second section]. Part 2. *Mem. Geol. Surv. Great Britain Palaeont.*, 4, 573–792.

DANZÉ, J. 1956. Contribution à l'étude des Sphenopterisé: les fougères sphénoptéridiennes du bassin houiller du Nord de la France. Études géologiques pour l'Atlas de topographie souterraine. I. Flore fossile.

DANZÉ-CORSIN, P. 1953. Contribution à l'étude des Marioptéridéesé: les Mariopteris du Nord de la France. Études géologiques pour l'Atlas de topographie souterraine. I. Flore fossile.

DAVIES, N. S., GARWOOD, R. J., MCMAHON, W. J., SCHNEIDER, J. W. and SHILLITO, A. P. 2021. The largest arthropod in Earth history: insights from newly discovered *Arthropleura* remains (Serpukhovian Stainmore Formation, Northumberland, England). *Journal of the Geological Society*, 179, jgs2021-115.

DERNOV, V. 2019. Taphonomy and paleoecology of fauna and flora from deltaic sandstones of Mospinka Formation (Middle Carboniferous) of Donets Basin. *GEO&BIO*, 2019, 37–63.

DOLLÉ, P. 1970. Sédimentation rythmée dans les charbons du Bassin Houiller du Nord-Pas de Calais.

GUTHÖRL, P. 1934. Die Arthropoden aus dem Carbon und Perm des -Saar-Nahe-Pfalz-Gebietes. *Abhandlungen der Preussischen Geologischen Landesanstalt, Neue Folge*, 164, 1–219.



——. 1935. Entdeckung und Bergung des größten, bis jetzt bekannten Exemplars von *Arthropleura armata* JORDAN & V. MEYER in Grube Maybach-Saar. *Zeitschrift der Deutschen Geologischen Gesellschaft*, 87, 687–692.

——. 1936. Arthropleura, der Riesengliederfuessler des Oberkarbons, und seine Verbreitung in den europaeischen Steinkohlenbecken. Berg- und Hüttenmännischen Zeitschrift, Glückauf", 72, 965–975.

HAHN, G., HAHN, R. and BRAUCKMANN, C. 1986. Zur Kenntnis von *Arthropleura* (Myriapoda; Ober-Karbon). *Geologica et Palaeontologica*, 20, 125–137.

HANNIBAL, J. T. 1997. Remains of *Arthropleura*, a gigantic myriapod arthropod, from the Pennsylvanian of Ohio and Pennsylvania. *Kirtlandia*, 50, 1–10.

HENNION, M., CUVELIER, J., GALLEGO, O. F., THIERRY, O. and VACHARD, D. 2015. Collections of the Natural History Museum of Lille. VIII. - Diplostraca (Types and figured specimens). *Annales de la Société Géologique du Nord*, 22 (2ème série), 109–118.

IZART, A., VASLET, D., BRIAND, C., BROUTIN, J., COQUEL, R., DAVY-DOV, V., DONSIMONI, M., EL WARTITI, M., ENSEPBAEV, T., GELUK, M., GOREVA, N., GÔRÜR, N., IQBAL, N., JOLTAEV, G., KOSSOYAVA, O. L., KRAINER, K., LAVEINE, J. P., MAKHLINA, M. K., MASLO, A., NEMIROVSKAYA, T., KORA, M., KOZITSKAYA, R., MASSA, D., MERCIER, D., MONOD, O., OPLUSTIL, S., SCHNEIDER, J. W., SCHÖNLAUB, H., STSCHEGOLEV, A., SÜSS, P., VACHARD, D., VAI, G. B., VOZÁROVÁ, A., WEISSBRODT, T. and ZDANOWSKI, A. 1998. Stratigraphic correlations between the continental and marine Tethyan and Peri-Tethyan basins during the Late Carboniferous and the Early Permian. *Geodiversitas*, 20, 521–595.

JORDAN, H. and MEYER, H. von. 1854. Ueber die Crustaceen der Steinkohlenformation von Saarbrücken: Faszinierende Einblicke in urzeitliche Fossilienwelten. *Palaeontographica*, 4, 1–15.

KRAUS, O. 2005. On the structure and biology of *Arthropleura* species (Atelocerata, Diplopoda; Upper Carboniferous/Lower Permian). *Verhandlungen des naturwissenschaftlichen Vereins Hamburg, Neue Folge*, 41, 5–23.

— and BRAUCKMANN, C. 2003. Fossil giants and surviving dwarfs. Arthropleurida and Pselaphognatha (Atelocerata, Diplopoda): characters, phylogenetic relationships and construction. *Verdhandlungen des Naturwissenschaftlichen Vereins in Hamburg*, 40, 5–50.

LAVEINE, J.-P. 1967. Contribution a l'étude de la flore du terrain houiller. Les Neuroptéridées du Nord de la France. Études géologiques pour l'Atlas de topographie souterraine. I. Flore fossile.

— and GOUBET, P. 1995. Mise au point sur Palaeoweichselia defrancei (ptéridospermée du Carbonifère), à partir d'un bel échantillon récolté dans le bassin houiller du nord de la France. Sciences Géologiques, bulletins et mémoires, 48, 21–47.

and BELHIS, A. 2007. Frond architecture of the seed-fern *Macroneuropteris scheuchzeri*, based on Pennsylvanian specimens from the Northern France coal field. *Palaeontographica B*, 277, 1–41.

——, COQUEL, R. and BROUSMICHE, C. 1978. L'espèce *Pecopteris simoni* (Zeiller, 1886–1888) sunonyme posterieur de *Palaeoweichselia defrancei* (Brongniart, 1828). *Geobios*, 11, 755–759.

LEMOIGNE, Y. and ZHANG, S. 1992. Pangaea, Palaeotethys and Paripteris. Comptes rendus de l'Académie des sciences. Série 2, Mécanique, Physique, Chimie, Sciences de l'univers, Sciences de la Terre, 314, 1103–1110.

—, —, LI, X., WU, X., ZHANG, S., ZHAO, X., ZHU, W. and ZHU, J. 1987. Paléogéographie de la Chine au Carbonifère à la lumière des données paléobotaniques, par comparaison avec les assemblages carbonifères d'Europe occidentale. *Comptes rendus de l'Académie des sciences. Série 2, Mécanique, Physique, Chimie, Sciences de l'univers, Sciences de la Terre*, 304, 391–394.

LE GALL, B. 1994. Deformation of the Nord-Pas-de-Calais Carboniferous Coalfield (France) in the Variscan Frontal Tectonic Pattern. In MASCLE, A. (ed.) *Hydrocarbon and Petroleum Geology of France. Special Publication of the European Association of Petroleum Geoscientists*, Vol. 4. Springer, Berlin, Heidelberg, 379-398 pp.

LHÉRITIER, M., EDGECOMBE, G. D., GARWOOD, R. J., BUISSON, A., GERBE, A., KOCH, N. M., VANNIER, J., ESCARGUEL, G., ADRIEN, J., FERNANDEZ, V., BERGERET-MEDINA, A. and PERRIER, V. 2024. Head anatomy and phylogenomics show the Carboniferous giant *Arthropleura* belonged to a millipede-centipede group. *Science Advances*, 10, eadp6362.

LUCAS, S. G., SCHNEIDER, J. W., NIKOLAEVA, S. and WANG, X. 2022. The Carboniferous timescale: an introduction. *Geological Society, London, Special Publications*, 512, 1–17.

MINELLI, A. 2011. Treatise on Zoology - Anatomy, Taxonomy, Biology. The Myriapoda. Vol. 1. Brill, Leidené; Boston.

——. 2015. Treatise on Zoology - Anatomy, Taxonomy, Biology. The Myriapoda. Vol. 2. Brill, Leiden, Boston.

MOLINA-SOLÍS, A., CLEAL, C. J., CAPEL, E. and CASCALES-MINANA, B. 2024. Vegetation dynamics in the Early–Middle Pennsylvanian coal swamps of the Nord-Pas-de-Calais Coalfield, France. *Palaeoworld*, 33, 328–340.

PACYNA, G., FLORJAN, S. and BORZECKI, R. 2012. New morphological features of *Arthropleura* sp. (Myriapoda, Diplopoda) based on new specimens from the upper Carboniferous of Lower Silesia (Poland). *Annales Societatis Geologorum Poloniae*, 82, 121–126.

PAPROTH, E., DUSAR, M., BLESS, M. J., BOUCKAERT, J., DELMER, A., FAIRON-DEMARET, M., HOULLEBERGHS, E., LALOUX, M., PIERART, P. and SOMERS, Y. 1983. Bio-and lithostratigraphic subdivisions of the Silesian in Belgium, a review. *Annales de la Société géologique de Belgique*, 106.

PAVELA, M. 2018. Fossil species from the Carboniferous: *Arthropleura* Jordan & Meyer 1854. *PALEO*, 74.

PILLOLA, G. L. and ZOBOLI, D. 2021. First occurrence of *Arthropleura armata* (Myriapoda) in the Moscovian (Carboniferous) of SW Sardinia (Italy). *Bollettino della Società Paleontologica Italiana*, 60, 1–6.

PROCTOR, C. J. 1998. Arthropleurids from the Westphalian D of Writhlington Geological Nature Reserve, Somerset. *Proceedings of the Geologists' Association*, 109, 93–98.

PRUVOST, P. 1912. Note sur un myriapode du terrain houiller du Nord. *Annales de la Societe Geologique du Nord*, 41, 63–68.

———. 1914. Nouvelles découvertes d'insectes fossiles dans le terrain houiller du Nord et du Pas-de-Calais. *Annales de la Société Géologique du Nord*, 43, 282–295.

———. 1919. Introduction à l'étude du terrain houiller du Nord et du Pasde-Calais: la faune continentale du terrain houiller du Nord de la France. Imprimerie nationale.

——. 1922. Les arachnides fossiles du Houiller de Belgique. Annales de la Société Scientifique de Bruxelles, 41, 349–355.

——. 1930. La faune continentale du terrain houiller de la Belgique. *Mémoires du Musée royal d'histoire naturelle de Belgique*, 103–282.

ŘEHOŘ, F. and ŘEHOŘOVA, M. 1972. Die Makrofauna des kohlenführenden Karbons im tschechoslovakischen Teil des oberschlesischen Beckens. *Pràci Venujeme 100. Výroči Ostravského Muzea*, 107–131.

ROLFE, W. I. 1985. Aspects of the Carboniferous terrestrial arthropod community. *9th Congress International Stratigraphy and Geology Carboniferous, Champaign-Urbana USA*, 5, 303–316.

RÖSSLER, R. and SCHNEIDER, J. W. 1997. Eine bemerkenswerte Paläobiocoenose im Unterkarbon Mitteleuropas–Fossilführung und Paläoenvironment der Hainichen-Subgruppe (Erzgebirge-Becken). Veröffentlichungen des Museums für Naturkunde Chemnitz, 20, 5–44.

——, ZIEROLD, T., FENG, Z., KRETZSCHMAR, R., MERBITZ, M., ANNACKER, V. and SCHNEIDER, J. W. 2012. A snapshot of an early Permian ecosystem preserved by explosive volcanism: new results from the Chemnitz petrified forest, Germany. *PALAIOS*, 27, 814–834.

SALTER, J. W. 1863. On some Species of Eurypterus and Allied Forms.



Quarterly Journal of the Geological Society of London, 19, 81–87.

SCHNEIDER, J. W., LUCAS, S. G., WERNEBURG, R. and RÖSSLER, R. 2010. Euramerican Late Pennsylvanian/early Permian arthropleurid/tetrapod associations - implications for the habitat and paleobiology of the largest terrestrial arthropod. *New Mexico Museum of Natural History and Science, Bulletin*, 49, 49–70.

SCHNEIDER, J. W., SCHOLZE, F., ROSS, A. J., BLAKE, B. M., Jr and LUCAS, S. G. 2022. Improved blattoid insect and conchostracan zonation for the Late Carboniferous, Pennsylvanian, of Euramerica. In LUCAS, S. G., SCHNEIDER, J. W., WANG, X. and NIKOLAEVA, S. (eds.) *The Carboniferous Timescale, Geological Society of London.* 

VALLOIS, B. and LEBRUN, P. 2010. Les plantes du Carbonifère supérieur de la concession houillère de Liévin (Pas-de-Calais). *Revue Française de Paléontologie Fossiles*, 2, 4–18.

— and NEL, A. 2023. Possible earliest evidence of insect pollination based on a new species of the Carboniferous medullosalean 'seed' genus *Pachytesta. Review of Palaeobotany and Palynology*, 316, 104948.

VERNON, R. D. 1912. On the Geology and Palæontology of the Warwickshire Coalfield. *Quarterly Journal of the Geological Society of London*, 68, 587–638.

WATERLOT, G. 1934. Bassin houiller de la Sarre et de la Lorraine. II. Faune fossile. Étude de la faune continentale du terrain houiller sarro-lorrain. *Études des Gîtes Minéraux de la France*, 25, 1–317.

WESENER, T., SIERWALD, P. and WÄGELE, J.-W. 2014. Sternites and spiracles – The unclear homology of ventral sclerites in the basal millipede order Glomeridesmida (Myriapoda, Diplopoda). *Arthropod Structure & Development*, 43, 87–95.

WILSON, H. M. 1999. Palaeobiology of the Arthropleuridea.Unpublished Ph.D., The University of Manchester, 312pp.

WOODWARD, H. 1907. V.—Further Notes on the Arthropoda of the British Coal-Measures. *Geological Magazine*, 4, 539–549.

ZEILLER, R. 1886. Bassin houiller de Valenciennes. Description de la flore fossile. Études des Gîtes Minéraux de la France. Pls I-XCIV. Paris.

———. 1894. Sur les subdivisions du Westphalien du Nord de la France d'après les caractères de la flore. *Bull. Soc. Géol. de France*, 3 ème Série; Tome XXII.

———. 1896. Etude sur quelques plantes fossiles, en particulier Vertebraria et Glossopteris, des environs de Johannesburg (Transvaal). *Bull. Soc. Géol. de France*, 24, 349–378.